



OFFICE OF THE PRINCIPAL
S.B.DEORAH COLLEGE
ULUBARI, GUWAHATI – 781 007
Tel.: 0361-3511878
E-mail: sbdeorahcollege@gmail.com
Website: www.sbdeorahcollege.org.in



The Institution has several collaborations/linkages for Faculty exchange, Student exchange, Internship, Field trip, On-the- job training, research etc. during the year

Research Collaboration

25. Dr. Jayantha K. Nath, Assistant Professor, Department of Chemistry, S. B. Deorah College, has Research Collaboration with Dr. Mrinmoy Manash Bharali, Professor, Dept. of Organic Synthesis and Process Chemistry, CSIR, Indian Institute of Chemical Technology and Academy of Scientific and Innovative Research, India.

Published on 12 April 2023. Downloaded by North East Institute of Science & Technology (RII) - Kirhaty (CSIR) on 4/20/2023 10:58:19 AM.

Green Chemistry

ROYAL SOCIETY OF CHEMISTRY

COMMUNICATION

View Article Online
View Journal

Check for updates

Utilization of methanol and ethanol for 3,3'-bis(indolyl)methane synthesis through activation of peroxymonosulfate over a copper catalyst†

Arpita Devi,^a Mrinmoy Manash Bharali,^{b,c} Subir Biswas,^a Tonmoy J. Bora,^a Jayanta K. Nath,^d Seonghwan Lee,^e Young-Bin Park,^e Lakshi Saikia,^f Manash J. Baruah^a and Kusum K. Bania^{g,h}

A greener and simple catalytic system is developed for the synthesis of biologically important 3,3'-bisindolyl(methanes) (BIMs) using C1 and C2 alcohols as the carbon source for the bridging methylene group. The reaction occurred under very mild and environment friendly conditions without the requirement of any toxic solvents. The low cost CuO-peroxymonosulfate (CuO-PMS) system allows the reaction to be highly efficient, resulting in very good product yield.

3,3'-Bisindolyl(methanes) (BIMs) are considered to be important compounds as they serve as structural motifs for many biologically active molecules.¹ In general, the Friedel-Crafts reaction of indoles with different types of aldehydes is mostly employed for the synthesis of such pharmaceutically important compounds.^{1a,c} However, the use of formaldehyde or acetaldehyde in the synthesis of 3,3'-BIMs is not considered to be an environmentally benign process as they are highly volatile and difficult to control under ambient conditions, and thereby cause damage to the atmosphere.^{1a} In order to overcome such problems, dehydrogenation of methanol (CH₃OH) and ethanol (C₂H₅OH) was employed as an alternative approach for the introduction of the bridging methylene group during the synthesis of 3,3'-BIMs.^{1c} The reported studies, however, used costly Ir or Ru based catalysts.^{1c-3} Apart from this, the reaction conditions required high temperature in the range of 120–150 °C, longer reaction time (18–20 h) and other additives such as an external base (Scheme 1a). Therefore, finding a suitable low cost metal catalyst or a catalytic system that can promote similar C–C bond formation reactions through the utilization of C1 and C2 alcohols in the formation of 3,3'-BIMs at a low temperature, with a short reaction time and with high selectivity would be highly beneficial (Scheme 1b).

In this regard, a CuO based catalyst is found to be highly effective for the dehydrogenation of CH₃OH or C₂H₅OH.⁴ Very recently, we have also demonstrated that a CuO catalyst loaded over magnesium-exchanged zeolite-Y (Mg²⁺-Y) can dehydrogenate C₂H₅OH and can be utilized for the synthesis of various cinnamaldehydes.⁵ At the same time, CuO is also known to activate peroxymonosulfate (PMS) which is used as an oxidant in the synthesis of C2/C3-trimerized indoles.⁶ However, no literature is available on the application of the CuO-PMS system

(a) Previous protocols:

(i) Ir/Mg₂AlO₄, 150 °C, 4 h^{1(a)}
(ii) Ru pincer complex, 135 °C, 18 h, Toluene, KOH, argon atmosphere^{1(b)}
(iii) [Cp*CrCl₂]₂, 150 °C, 12 h, KOtBu^{1(c)}

(b) Present Protocol:

CuO-PMS
RT, 20 min

11 examples (upto 85% yield)

PMS= Peroxymonosulfate
RT= Room temperature

- Low cost and recyclable catalyst
- Ambient reaction conditions
- Less duration

Scheme 1 Approaches for the synthesis of 3,3'-BIMs using methanol or ethanol as the source of the methylene group.