

Total number of printed pages-11

**3 (Sem-5/CBCS) MAT HC 2**

**2022**

**MATHEMATICS**

(Honours)

Paper : MAT-HC-5026

**(Linear Algebra)**

Full Marks : 80

Time : Three hours

**The figures in the margin indicate full marks for the questions.**

1. Answer **any ten** questions :  $1 \times 10 = 10$

(i) "A plane in  $\mathbb{R}^3$  not through the origin is a subspace of  $\mathbb{R}^3$ ."

(State True or False)

(ii) If the equation  $AX = 0$  has only the trivial solution then what is the null space of  $A$ ?

(iii) Suppose two matrices are row equivalent. Are their row spaces the same?

Contd.

(iv) Let  $A$  be matrix of order  $m \times n$ . When the column space of  $A$  and  $\mathbb{R}^m$  are equal?

(v) Is the set  $\{\sin t, \cos t\}$  linearly independent in  $C[0, 1]$ ?

(vi) What is the dimension of zero vector space?

(vii) If  $A$  is a  $7 \times 9$  matrix with a two-dimensional null space, what is the rank of  $A$ ?

(viii) "0 is an eigenvalue of a matrix  $A$  if and only if  $A$  is invertible."

(State True **or** False)

(ix) Let  $A$  be an  $n \times n$  matrix such that determinant of  $A$  is zero. Is  $A$  invertible?

(x) When two matrices  $A$  and  $B$  are said to be similar?

(xi) Define complex eigenvalue of a matrix.

(xii) Let an  $n \times n$  matrix has  $n$  distinct eigenvalues. Is it diagonalizable?

(xiii) What do you mean by distance between two vectors in  $\mathbb{R}^n$ ?

(xiv) Which vector is orthogonal to every vector in  $\mathbb{R}^n$ ?

(xv) Is inner product of two vector  $u$  and  $v$  in  $\mathbb{R}^n$  commutative?

(xvi) "An orthogonal matrix is invertible."  
(State True or False)

(xvii) If the number of free variables in the equation  $Ax = 0$  is  $p$ , then what is the dimension of null space of  $A$ ?

(xviii) Let  $T$  be a linear operator on a vector space  $V$ . Is the subspace of  $\{0\}$  of  $V$   $T$ -invariant?

2. Answer **any five** questions :  $2 \times 5 = 10$

(i) Show that the set  $H$  of all points of  $\mathbb{R}^2$  of the form  $(3r, 2 + 5r)$  is not a vector space.

(ii) Let  $A = \begin{bmatrix} 2 & 4 & -2 & 1 \\ -2 & -5 & 7 & 3 \\ 3 & 7 & -8 & 6 \end{bmatrix}$  and let

$u = \begin{bmatrix} 3 \\ -2 \\ -1 \\ 0 \end{bmatrix}$ . Is  $u$  in null space of  $A$ ?

(iii) In  $\mathbb{R}^3$ , show that the set  $W = \{(a, b, c) : a^2 + b^2 + c^2 \leq 1\}$  is not a subset of  $V$ .

(iv) Let  $P_1(t) = 1, P_2(t) = t, P_3(t) = 4 - t$ . Show that  $\{P_1, P_2, P_3\}$  is linearly dependent in the vector space of polynomials.

(v) Let  $A = \begin{bmatrix} 1 & 6 \\ 5 & 2 \end{bmatrix}, u = \begin{bmatrix} 6 \\ -5 \end{bmatrix}$ . Examine whether  $u$  is an eigenvector of  $A$ .

(vi) The characteristic polynomial of a  $6 \times 6$  matrix is  $\lambda^6 - 4\lambda^5 - 12\lambda^4$ . Find the eigenvalue of the matrix.

(vii) Show that the eigenvalues of a triangular matrix are just the diagonal elements of the matrix.

(viii) Let  $v = (1, -2, 2, 0)$ . Find a unit vector  $u$  in the same direction as  $v$ .

(ix) Let  $u = \begin{bmatrix} -2 \\ 1 \end{bmatrix}$  and  $v = \begin{bmatrix} -3 \\ 1 \end{bmatrix}$ . Compute  $\frac{u \cdot v}{u \cdot u}$ .

$\lambda^2(\lambda-6)^2 + 2(\lambda-6)$   
 $\lambda^2(\lambda-6)^2 + 2(\lambda-6)$   
 $\lambda^2(\lambda-6)^2 + 2(\lambda-6)$

(x) Suppose  $S = \{u_1, u_2, \dots, u_n\}$  contains a dependent subset. Show that  $S$  is also dependent.

3. Answer **any four** questions :  $5 \times 4 = 20$

(i) Let  $A = \begin{bmatrix} 2 & 4 & -2 & 1 \\ -2 & -5 & 7 & 3 \\ 3 & 7 & -8 & 6 \end{bmatrix}$ . Find a non-

zero vector in column space of  $A$  and a non-zero vector in null space of  $A$ .

(ii) If a vector space  $V$  has a basis  $B = \{b_1, b_2, \dots, b_n\}$ , then prove that any set in  $V$  containing more than  $n$  vectors must be linearly dependent.

(iii) Let  $B = \{b_1, b_2, \dots, b_n\}$  be a basis for a vector space  $V$ , then prove that the co-ordinate mapping  $x \rightarrow [x]_B$  is a one-to-one linear transformation from  $V$  onto  $\mathbb{R}^n$ .

(iv) Prove that similar matrices have the same characteristic polynomial and hence the same eigenvalues.

(v) Is 5 an eigenvalue of  $A = \begin{bmatrix} 6 & -3 & 1 \\ 3 & 0 & 5 \\ 2 & 2 & 6 \end{bmatrix}$ ?

(vi) Let  $U = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{2}{3} \\ \frac{1}{\sqrt{2}} & -\frac{2}{3} \\ 0 & \frac{1}{3} \end{bmatrix}$  and  $x = \begin{bmatrix} \sqrt{2} \\ 3 \end{bmatrix}$ . Show

that  $U$  has orthonormal columns and  $\|Ux\| = \|x\|$ .

(vii) Find a QR factorization of

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}.$$

(viii) Find the range and kernel of

$$T: \mathbb{R}^2 \rightarrow \mathbb{R}^2 \text{ defined by } \begin{bmatrix} x \\ y \end{bmatrix} \rightarrow \begin{bmatrix} x+y \\ x-y \end{bmatrix}.$$

4. Answer **any four** questions:  $10 \times 4 = 40$

(i) Find the spanning set for the null space of the matrix

$$A = \begin{bmatrix} -3 & 6 & -1 & 1 & -7 \\ 1 & -2 & 2 & 3 & -1 \\ 2 & -4 & 5 & 8 & -1 \end{bmatrix}.$$

(ii) Let  $S = \{v_1, v_2, \dots, v_r\}$  be a set in a vector space  $V$  over  $\mathbb{R}$  and let  $H = \text{span}\{v_1, v_2, \dots, v_r\}$ . Prove that—

(a) if one of the vectors in  $S$  is a linear combination of the remaining vectors in  $S$ , then the set formed from  $S$  by removing that vector still spans  $H$ ;

(b) if  $H \neq \{0\}$ , some subset of  $S$  is a basis for  $H$ .

5+5=10

(iii) Let  $V$  be the vector space of  $2 \times 2$  symmetric matrices over  $\mathbb{R}$ . Show that  $\dim V = 3$ . Also find the co-ordinate vector of the matrix

$$A = \begin{bmatrix} 4 & -11 \\ -11 & -7 \end{bmatrix} \text{ relative to the basis}$$

$$\left\{ \begin{bmatrix} 1 & -2 \\ -2 & 1 \end{bmatrix}, \begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix}, \begin{bmatrix} 4 & -1 \\ -1 & -5 \end{bmatrix} \right\}.$$

5+5=10

(iv) Define a diagonalizable matrix. Prove that an  $n \times n$  matrix  $A$  is diagonalizable if and only if  $A$  has  $n$  linearly independent eigenvectors. 1+9=10

(v) (a) Show that  $\lambda$  is an eigenvalue of an invertible matrix  $A$  if and only if  $\lambda^{-1}$  is an eigenvalue of  $A^{-1}$ .

(b) If  $\lambda_1, \lambda_2, \dots, \lambda_n$  are the eigenvalues of  $A$ , then show that  $k\lambda_1, k\lambda_2, \dots, k\lambda_n$  are the eigenvalues of  $kA$ .

(c) Show that the matrices  $A$  and  $A^T$  (transpose of  $A$ ) have the same eigenvalues.

5+2½+2½=10



(vi) Compute  $A^8$  where  $A = \begin{bmatrix} 4 & -3 \\ 2 & -1 \end{bmatrix}$ .

(vii) Define orthogonal set and orthogonal basis of  $\mathbb{R}^n$ . Show that  $S = \{u_1, u_2, u_3\}$  is an orthogonal basis for  $\mathbb{R}^3$ . Also

express the vector  $y = \begin{bmatrix} 6 \\ 1 \\ -8 \end{bmatrix}$  as a linear

combination of the vector in  $S$ .

$$(1+1)+5+3=10$$

(viii) Let  $V$  be an inner product space. Show that—

(a)  $\langle v, 0 \rangle = \langle 0, v \rangle = 0$ ;

(b)  $\langle u, v+w \rangle = \langle u, v \rangle + \langle u, w \rangle$

where  $u, v, w \in V$ ;

(c) Define norm of a vector in  $V$ ;

(d) For  $u, v$  in  $V$ , show that

$$|\langle u, v \rangle| \leq \|u\| \|v\|.$$

$$2+2+1+5=10$$

(ix) What do you mean by Gram-Schmidt process? Prove that if  $\{x_1, x_2, \dots, x_p\}$

is a basis for a subspace  $W$  of  $\mathbb{R}^n$  and define  $v_1 = x_1$

$$v_2 = x_2 - \frac{x_2 \cdot v_1}{v_1 \cdot v_1} v_1$$

$$v_3 = x_3 - \frac{x_3 \cdot v_1}{v_1 \cdot v_1} v_1 - \frac{x_3 \cdot v_2}{v_2 \cdot v_2} v_2$$

$$v_p = x_p - \frac{x_p \cdot v_1}{v_1 \cdot v_1} v_1 - \frac{x_p \cdot v_2}{v_2 \cdot v_2} v_2 - \dots - \frac{x_p \cdot v_{p-1}}{v_{p-1} \cdot v_{p-1}} v_{p-1}$$

then  $\{v_1, v_2, \dots, v_p\}$  is an orthogonal basis for  $W$ . Also if  $W = \text{span}\{x_1, x_2\}$

where  $x_1 = \begin{bmatrix} 3 \\ 6 \\ 6 \end{bmatrix}$ ,  $x_2 = \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}$ . Construct an

orthogonal basis  $\{v_1, v_2\}$  for  $W$ .

$$1+6+3=10$$

(x) Define orthogonal complement of a subspace. Let  $\{u_1, u_2, \dots, u_5\}$  be an orthogonal basis for  $\mathbb{R}^5$  and  $y = c_1u_1 + \dots + c_5u_5$ . If the subspace  $W = \text{span}\{u_1, u_2\}$  then write  $y$  as the sum of vectors  $Z_1$  in  $W$  and a vector  $Z_2$  in complement of  $W$ . Also find the distance from  $y$  to  $W = \text{span}\{u_1, u_2\}$ ,

$$\text{where } y = \begin{bmatrix} -1 \\ -5 \\ 10 \end{bmatrix}, u_1 = \begin{bmatrix} 5 \\ -2 \\ 1 \end{bmatrix}, u_2 = \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}.$$

$$1+6+3=10$$

---