

Total number of printed pages-12

1 (Sem-4) PHY 1

2025

PHYSICS

Paper : PHY0400104

(Classical Mechanics)

Full Marks : 60

Time : 2½ hours

The figures in the margin indicate full marks for the questions.

1. Answer the following questions : $1 \times 8 = 8$

(a) How many degrees of freedom are possessed by a ball moving on the surface of a sphere ?

(b) Lagrangian of a free particle moving along X-axis is given by $L = \frac{1}{2}m\dot{x}^2$.

What is its generalised momentum ?

(c) Which one of the following is a correct expression for Legendre transformation ?

(i) $H = \sum \dot{p}_j q_j - L$

(ii) $H = \sum p_j \dot{q}_j + L$

(iii) $H = \sum p_j \dot{q}_j - L$

(iv) $H = \sum \dot{p}_j \dot{q}_j - L$

(d) Lagrangian of a particle moving in a central force potential $V(r)$ is expressed as—

$$L = \frac{1}{2} m \dot{r}^2 + \frac{1}{2} m r^2 \dot{\theta}^2 + \frac{1}{2} m r^2 \sin^2 \theta \dot{\phi}^2 - V(r).$$

Which one of the following is a correct statement ?

- (i) Momentum conjugate to γ is conserved.
- (ii) Momentum conjugate to θ is conserved.
- (iii) Momentum conjugate to ϕ is conserved.
- (iv) Energy is not conserved.

(e) If $V(x)$ is potential energy of a particle moving along x -direction which one of the following is a condition of stable equilibrium ?

(i) $V(x) = 0, \frac{dV}{dx} = 0$

(ii) $\frac{dV}{dx} = 0, \frac{d^2V}{dx^2} < 0$

(iii) $\frac{dV}{dx} = 0, \frac{d^2V}{dx^2} > 0$

(iv) $\frac{dV}{dx} \rightarrow \infty, \frac{d^2V}{dx^2} > 0$

(f) Which one of the following is a correct statement in special relativity ?

- (i) Velocity of light depends on velocities of the observers.
- (ii) If two events are simultaneous in one frame they are simultaneous in all other frames.
- (iii) If two events are simultaneous in one frame they are not simultaneous other frames.
- (iv) Mass of a body reduces to zero when its velocity approaches velocity of light.

(g) If momentum of a particle is $p = 2mc$, which one of the following is the correct expression for energy of the particle as per relativistic energy momentum relation ?

(i) $E = \pm 5mc^2$

(ii) $E = \pm \sqrt{5}mc^2$

(iii) $E = \pm 4mc^2$

(iv) $E = \pm 2mc^2$

(h) If \vec{u} is velocity of a fluid element, which one of the following represents as incompressible fluid ?

(i) $\nabla^2 \vec{u} = 0$

(ii) $(\vec{u} \cdot \vec{\nabla}) \vec{u} = 0$

(iii) $\vec{\nabla} \cdot \vec{u} = 0$

(iv) $\vec{\nabla} u^2 = 0$

2. Answer **any six** questions : 2×6=12

(a) Lagrangian of a simple pendulum of unit mass is given by

$$L = \frac{1}{2} l^2 \dot{\theta}^2 - gl(1 - \cos \theta).$$

Obtain the Euler-Lagrange equation.

(b) Lagrangian of a particle moving along X-direction is

$$L = \frac{1}{2} m \dot{x}^2 - \frac{1}{2} k x^4.$$

Obtain the Hamiltonian of the particle.

(c) In spherical polar coordinates Lagrangian of a free particle is given by

$$L = \frac{1}{2} m \dot{r}^2 + \frac{1}{2} m r^2 \dot{\theta}^2 + \frac{1}{2} m r^2 \sin^2 \theta \dot{\phi}^2.$$

Obtain the generalised momentum conjugate to ϕ when the particle moves

in equatorial plane $\theta = \frac{\pi}{2}$.

- (d) Lagrangian of a particle attached to a spring of spring constant K is

$$L = \frac{1}{2}m\dot{x}^2 - \frac{1}{2}kx^2. \text{ Reduce Hamilton's}$$

Canonical equation $\dot{p}_x = -\frac{\partial H}{\partial x}$ in this case to the following form $m\ddot{x} = -kx$.

- (e) A particle is displaced by an amount $x - x_0$ from its equilibrium position $x = x_0$. Obtain the Taylor expansion of potential energy $V(x)$ around the equilibrium $x = x_0$.

- (f) Write down the *two* postulates of special relativity.

- (g) Lorentz transformation for time is given by

$$t' = \gamma \left(t - \frac{vx}{C^2} \right). \text{ Show that if two events}$$

are simultaneous in one frame they are not simultaneous in the other frame.

- (h) Calculate the energy equivalent to mass of the Sun, $M = 2 \times 10^{30} \text{ kg}$.

- (i) Show that time derivative of velocity (\bar{u}) of a fluid element is

$$\frac{d\bar{u}}{dt} = \frac{\partial \bar{u}}{\partial t} + (\bar{u} \cdot \bar{\nabla}) \bar{u}.$$

- (j) What is an ideal fluid? Write down the equation of continuity.

3. Answer **any four** questions : $5 \times 4 = 20$

- (a) What do you mean by stable equilibrium? If $q_i = q_{0i} = \eta_i$ represents displacement of generalised coordinate from equilibrium (q_{0i}) expand the potential energy $V(q_1, q_2, \dots, q_n)$ in a Taylor series about q_{0i} and obtain the potential energy matrix V_{ij} ... writing the

kinetic energy as $T = \frac{1}{2} m_{ij} \dot{\eta}_i \dot{\eta}_j$ and

expanding the function m_{ij} in a Taylor series around q_{0i} obtain an appropriate expression for kinetic energy matrix.

1+2+2=5

(b) For a system in equilibrium derive the principle of virtual work. Apply appropriate assumption to obtain D' Alembert's principle. $2\frac{1}{2}+2\frac{1}{2}=5$

(c) Lagrangian for a simple pendulum is given by $L = \frac{1}{2}ml^2\dot{\theta}^2 - mgl(1 - \cos\theta)$.

Obtain the Hamiltonian and hence obtain Hamilton's Canonical equations. $3+2=5$

(d) Lagrangian of a particle in cylindrical coordinate system with potential energy $V(r, \theta, z)$ is given by

$$L = \frac{1}{2}m(\dot{r}^2 + r^2\dot{\theta}^2 + \dot{z}^2) - V(r, \theta, z)$$

obtain Euler-Lagrange equations for r , θ and z .

(e) Potential energy of a particle moving along X-axis is given by

$$V(x) = -\frac{1}{2}kx^2 + \lambda x^4 \quad (k, \lambda > 0).$$

Show that $x = 0, +\sqrt{\frac{k}{4\lambda}}$ and $-\sqrt{\frac{k}{4\lambda}}$ are equilibrium positions. Out of these three, identify the stable equilibrium positions. $2+3=5$

(f) What is the inadequacy of Galilean transformation? Derive length contraction and time dilation formulae from Lorentz transformation equations. $1+2+2=5$

(g) From Lorentz transformation equations of (x, t) obtain the relativistic velocity addition formula. Show that velocity of light is invariant. $4+1=5$

(h) If relativistic energy and momentum are written as

$$E = \frac{mc^2}{\sqrt{1-v^2/c^2}} \quad \text{and} \quad p = \frac{mv}{\sqrt{1-v^2/c^2}}$$

show that $\frac{E^2}{c^2} - p^2 = m^2c^2$.

Two particles, each of mass m collide head on at the speed of $V = \frac{3}{5}C$. They form a composite particle of mass M which is at rest. Use conservation of relativistic energy to show that

$$M = \frac{5}{2}m. \quad 3+2=5$$

4. Answer **any two** questions : $10 \times 2 = 20$

(a) Lagrangian for a particle moving under a central force potential $V(r)$ is expressed as

$$L = \frac{1}{2}m(\dot{r}^2 + r^2\dot{\theta}^2) - V(r).$$

Use Euler-Lagrange equation for θ to show that $P_\theta = mr^2\dot{\theta}$ is a conserved momentum. Show that a real velocity of the particle remains constant. Show that Euler-Lagrange equation for the coordinate r is

$$m\ddot{r} - mr\dot{\theta}^2 = f(r), \text{ where}$$

$$f(r) = -\frac{\partial V(r)}{\partial r}. \text{ Obtain Hamiltonian of}$$

the particle. Show that radial velocity of the particle is

$$\frac{dr}{dt} = \sqrt{\frac{2}{m} \left(E - V(r) - \frac{P_\theta^2}{2mr^2} \right)}.$$

$$2+2+2+2+2=10$$

(b) Show that Euler-Lagrange equation can be written as

$\dot{p}_i = \partial L / \partial q_i$, where p_i is the generalised momentum. If the Lagrangian is expressed as $L(q_i, \dot{q}_i, t)$ and Legendre transformation is given by

$H(q_i, p_i, t) = p_i \dot{q}_i - L(q_i, \dot{q}_i, t)$. Obtain Hamilton's Canonical equations.

$$2+8=10$$

(c) Write down Newton's second law of motion for a system of particles acted by external and internal forces. Define holonomic and non-holonomic constraints with equations and examples. A particle of mass m is falling freely under gravity vertically along Z -axis. Construct the Lagrangian. Obtain Hamilton's Canonical equation for the particle.

$$2+2+2+2+2=10$$

- (d) Mass of a relativistic particle changes with velocity as

$$m = \frac{m_0}{\sqrt{1 - v^2/C^2}}, \text{ where } m_0 \text{ is the}$$

rest mass. If velocity of the particle increases from 0 to v use work energy theorem to show that gain in kinetic energy of the particle is

$E_k = (m - m_0)C^2$. From this show that total relativistic energy of the particle

is $E = \frac{m_0 C^2}{\sqrt{1 - v^2/C^2}}$. 8+2=10

- (e) Show that Lorentz transformation reduces to Galilean transformation if $v \ll C$. Represent Lorentz transformation as rotation in spacetime. From Lorentz transformation equations for (x, y, z, t) , show that

$$c^2 t'^2 - x'^2 - y'^2 - z'^2 = c^2 t^2 - x^2 - y^2 - z^2.$$

2+5+3=10